

# Injective spaces and the filter monad

Martín Hötzel Escardó

Department of Computing, Edinburgh University

Version of Monday 3<sup>rd</sup> November 1997

## Abstract

An injective space is a topological space with a strong extension property for continuous maps with values on it. A certain filter space construction embeds every  $T_0$  topological space into an injective space. The construction gives rise to a monad. We show that the monad is of the Kock-Zöberlein type and apply this to obtain a simple proof of the fact that the algebras are the continuous lattices (Alan Day, 1975). In previous work we established an injectivity theorem for monads of this type, which characterizes the injective objects over a certain class of embeddings as the algebras. For the filter monad, the class turns out to consist precisely of the subspace embeddings. We thus obtain as a corollary that the injective spaces over subspace embeddings are the continuous lattices endowed with the Scott topology (Dana Scott, 1972). Similar results are obtained for continuous Scott domains, which are characterized as the injective spaces over *dense* subspace embeddings.

**Key-words:** Extension of maps, injective space, continuous lattice, continuous Scott domain, domain theory, Kock-Zöberlein monad.

**AMS classification:** 54C20, 54C25, 54C15, 54E99, 06B35, 18C20.

## 1 Introduction

An injective space is a topological space  $D$  with a strong extension property for continuous functions with values on  $D$ , to the extent that every continuous map  $f : X \rightarrow D$  extends to a continuous map  $\bar{f} : Y \rightarrow D$ , for every space  $Y$  containing  $X$  as a subspace. For example, the extended real line endowed with the topology of lower semicontinuity is an injective space.

A certain filter space construction embeds every  $T_0$  topological space into an injective space. In addition to contain the given space embedded as a subspace, the filter space appears as a quotient of its own filter space. Technically and more precisely, the filter space construction is a monad [7]. We show that the monad

is of the Kock-Zöberlein type [6] and apply this to obtain a simple proof of Alan Day's result that its algebras are the continuous lattices [2].

In [3] we proved that, given a category with the structure of a Kock-Zöberlein monad, the injective objects over a certain class of embeddings specified in term of the monad structure coincide with the algebras of the monad. It turns out that for the filter monad the embeddings are exactly the subspace embeddings. The fact that the injective spaces over subspace embeddings are precisely the continuous lattices endowed with the Scott topology, established by Dana Scott [8], thus appears as a corollary of the characterization of the algebras.

The continuous Scott domains are the algebras of the *proper* filter monad [11]. For this monad, the associated embeddings are precisely the *dense* subspace embeddings, and hence the injective spaces over dense embeddings are characterized as the continuous Scott domains. This characterization is folklore for the experts. It was first formulated and proved by Scott, but only published as Exercise II.3.19 of the *Compendium* [5], whose emphasis is on continuous lattices (personal communication).

More examples of the above situation, including the lower and upper space monads, are investigated in the paper [3], which also includes a long introduction explaining the fundamental role of injectivity in Scott's mathematical theory of computation and its connections with function spaces. Even more examples have been recently discovered by Bob Flagg and the author [4].

For background on continuous lattices the reader is referred to [5]. For more about domain theory and topology see [1, 9].

## 2 The filter space construction

The filter monad is defined on the category of  $T_0$  topological spaces and continuous maps. Recall that a space is  $T_0$  if no two distinct points share the same system of neighborhoods. Given a space  $X$ , one denotes its lattice of opens sets by  $\Omega X$  and constructs the **filter space**  $\mathcal{F}X$  as follows. The points are the filters of  $\Omega X$ . The open sets are generated by the sets

$$\square U = \{\phi \in \mathcal{F}X \mid U \in \phi\}, \quad U \in \Omega X,$$

which form a base as  $\square U \cap \square U' = \square(U \cap U')$ .

### 2.1 $\mathcal{F}X$ is an algebraic lattice endowed with the Scott topology.

It is immediate that the specialization order of  $\mathcal{F}X$  is inclusion of filters. Since the lattice of filters is algebraic with the principal filters as the compact elements, and since  $\square U$  is the set of filters containing the principal filter  $\uparrow U$ , we see that  $\square U$  is a basic Scott open set, which establishes the claim.

Given a map  $f : X \rightarrow Y$ , one defines a function  $\mathcal{F}f : \mathcal{F}X \rightarrow \mathcal{F}Y$  by

$$\mathcal{F}f(\phi) = \{V \in \Omega Y \mid f^{-1}(V) \in \phi\}.$$

Then  $\mathcal{F}$  is a functor and one has natural transformations  $\eta_X : X \rightarrow \mathcal{F}X$  and  $\mu_X : \mathcal{F}\mathcal{F}X \rightarrow \mathcal{F}X$  defined by

$$\eta_X(x) = \{U \in \Omega X \mid x \in U\}, \quad \mu_X(\Phi) = \bigcup \{\bigcap \mathcal{U} \mid \mathcal{U} \in \Phi\}.$$

which make  $\mathcal{F}$  into a monad  $\mathcal{F} = (\mathcal{F}, \eta, \mu)$ . Continuity of the functions defined above follows from the fact that

$$(\mathcal{F}f)^{-1}(\square V) = \square f^{-1}(V), \quad \eta_X^{-1}(\square U) = U, \quad \mu_X^{-1}(\square U) = \square \square U.$$

The first two equations are routinely verified and the last follows from §2.2 below. The second equation together with the fact that  $\eta_X$  is one-to-one shows that  $\eta_X$  embeds  $X$  as a subspace of  $\mathcal{F}X$ . The unit laws  $\mu_X \circ \eta_{\mathcal{F}X} = \mu_X \circ \mathcal{F}\eta_X = \text{id}_{\mathcal{F}X}$  show that  $\mu_X$  is a retraction map (in two ways) and hence  $\mathcal{F}X$  is a quotient of  $\mathcal{F}\mathcal{F}X$ , because retractions are quotient maps.

## 2.2 $\mu_X(\Phi) = \{U \in \Omega X \mid \square U \in \Phi\}$ .

Let  $U \in \mu_X(\Phi)$ . Then there is  $\mathcal{U} \in \Phi$  with  $U \in \bigcap \mathcal{U}$ , and hence with  $U \in \phi$  for all  $\phi \in \mathcal{U}$ . From this we see that  $\mathcal{U} \subseteq \square U$ . Hence  $\square U \in \Phi$  because filters are upper closed. Therefore  $\mu_X(\Phi) \subseteq \{U \in \Omega X \mid \square U \in \Phi\}$ . In order to establish the inclusion in the other direction, let  $U \in \Omega X$  with  $\square U \in \Phi$ . We have to show that there is  $\mathcal{U} \in \Phi$  with  $U \in \bigcap \mathcal{U}$ . We can take  $\mathcal{U} = \square U$ , because  $\bigcap \square U = \uparrow U$ , the principal filter generated by  $U$ .

## 3 Kock-Zöberlein monads

A monad  $T = (T, \eta, \mu)$  defined on a poset-enriched category  $\mathcal{X}$ , with  $T : \mathcal{X} \rightarrow \mathcal{X}$  a locally monotone functor, is said to be of the **Kock-Zöberlein type** if the condition

$$\eta_{TX} \sqsubseteq T\eta_X$$

holds for all  $X$ . Notice that our definition is dual (at the level of hom-posets) to that of [6].

Since  $\mathcal{X}$  is poset-enriched, one can consider adjunctions of arrows. Given arrows  $l : X \rightarrow Y$  and  $r : Y \rightarrow X$  one defines

$$l \dashv r \text{ iff } l \circ r \sqsubseteq \text{id}_Y \text{ and } \text{id}_X \sqsubseteq r \circ l,$$

and one says that  $l$  is **left adjoint** to  $r$  and that  $r$  is **right adjoint** to  $l$ . In this case each adjunct  $l$  and  $r$  is uniquely determined by the other. The adjunction is said to be **reflective** if  $l \circ r = \text{id}_Y$ , and **coreflective** if  $\text{id}_X = r \circ l$ . In these cases one writes  $l \dashv_r r$  and  $l \dashv_c r$  respectively.

By specializing Anders Kock's results [6] from 2-categories to poset-enriched categories, we learn that

**3.1** If  $T = (T, \eta, \mu)$  is a Kock-Zöberlein monad then

1. An arrow  $\alpha : TX \rightarrow X$  is the structure map of a  $T$ -algebra iff  $\eta_X \dashv_c \alpha$ .
2.  $\eta_{TX} \dashv \mu_X$ .
3.  $\mu_X \dashv T\eta_X$ .

(We showed in [3] that each of these properties is in fact equivalent to the Kock-Zöberlein property.)

By §3.1(1), every object can be the underlying object of at most one algebra, and every structure map of an algebra is uniquely determined by the underlying object of the algebra (as the right adjoint of the unit of the object). Due to this reason we can identify the algebras of a Kock-Zöberlein monad with their underlying objects.

## 4 Injective objects which are the algebras of Kock-Zöberlein monads

In what follows we work with a Kock-Zöberlein monad  $T = (T, \eta, \mu)$  defined on a poset-enriched category  $\mathcal{X}$ . The arrows singled out in the following definition are particular cases of the *semiupper* maps of [10], for which the reflectivity condition is not required:

**4.1** By a  *$T$ -embedding* we mean an arrow  $j : X \rightarrow Y$  such that the map  $Tj : TX \rightarrow TY$  has a reflective left adjoint, denoted by  $T^*j : TY \rightarrow TX$ .

For example,  $\eta_X : X \rightarrow TX$  is a  $T$ -embedding with  $T^*\eta_X = \mu_X$ , because the adjunction 3.1(3) is reflective by virtue of the unit law  $\mu_X \circ T\eta_X = \text{id}_X$ .

**Note** *The following conditions are equivalent:*

1.  $T$ -embeddings are order-monic.
2. Each component of  $\eta$  is order-monic.
3.  $T$  is order-faithful.

(1)  $\implies$  (2): Immediate. (2)  $\implies$  (3): If  $Tf \sqsubseteq Tg$  then we have that  $Tf \circ \eta_X \sqsubseteq Tg \circ \eta_X$  by composition with  $\eta_X$ , that  $\eta_Y \circ f \sqsubseteq \eta_Y \circ g$  by naturality, and that  $f \sqsubseteq g$  by the assumption. (3)  $\implies$  (1): Let  $j : X \rightarrow Y$  be a  $T$ -embedding and  $f, g : Z \rightarrow X$  be arrows with  $j \circ f \sqsubseteq j \circ g$ . Then  $Tj \circ Tf \sqsubseteq Tj \circ Tg$  by local monotonicity of  $T$ . Hence  $Tf \sqsubseteq Tg$  because  $Tj$  is split-mono. Therefore  $f \sqsubseteq g$  by the assumption, and the proof is concluded.

An object  $D$  is said to be **injective** over a class of arrows  $J$  if every map  $f : X \rightarrow D$  has an extension  $\bar{f} : Y \rightarrow D$  along any  $j : X \rightarrow Y \in J$ , in the sense that the following diagram commutes:

$$\begin{array}{ccc} X & \xrightarrow{j} & Y \\ & \searrow f & \nearrow \bar{f} \\ & & D \end{array}$$

One normally assumes that  $J$  consists of monomorphisms, so that the word *extension* is applied in the usual sense, but this is unimportant here. Notice that the extension  $\bar{f}$  need not be unique. But since  $\mathcal{X}$  is poset-enriched, a definition with canonical choice is possible. We first recall a concept.

A **right Kan extension** of a map  $f : X \rightarrow D$  along an arrow  $j : X \rightarrow Y$  is a (necessarily unique) arrow  $f/j : Y \rightarrow D$  such that

1.  $f/j \circ j \sqsubseteq f$  and
2.  $\bar{f} \circ j \sqsubseteq f$  implies  $\bar{f} \sqsubseteq f/j$ .

In other words,  $f/j$  is the largest solution in  $\bar{f}$  to the inequality  $\bar{f} \circ j \sqsubseteq f$ . In the case that we have equality in (1), so that  $f/j$  is not only a right Kan extension but also an actual extension in the above liberal sense, we say that  $f/j$  is a **right extension** of  $f$  along  $j$ .

**4.2** We say that an object  $D$  is **right injective** over  $J$  if every  $f : X \rightarrow D$  has a right extension  $f/j : Y \rightarrow D$  along any  $j : X \rightarrow Y \in J$ .

**4.3** The following statements are equivalent for any object  $D$ :

1.  $D$  is injective over  $T$ -embeddings.
2.  $D$  is right injective over  $T$ -embeddings.
3.  $D$  is a  $T$ -algebra.

In this case, if  $f : X \rightarrow D$  is any arrow and  $j : X \rightarrow Y$  is a  $T$ -embedding then  $f/j = m_D \circ Tf \circ T^*j \circ \eta_Y$ .

Here  $m_D : TD \rightarrow D$  is the unique structure map of the algebra  $D$ . The construction of  $f/j$  is illustrated in the following diagrams:

$$\begin{array}{ccc} X & \xrightarrow{j} & Y \\ & \searrow f & \nearrow f/j \\ & & D \end{array} \qquad \begin{array}{ccccc} TX & \xleftarrow{T^*j} & TY & \xleftarrow{\eta_Y} & Y \\ & \searrow Tf & & & \\ & & TD & & \\ & & \searrow m_D & & \\ & & & & D \end{array}$$

This was established in [3].

**Note** It is a basic property of monads that  $TX$  endowed with the structure map  $\mu_X : TTX \rightarrow TX$  is a free  $T$ -algebra with insertion of generators  $\eta_X : X \rightarrow TX$ . More precisely, the following universal property holds. Given any  $T$ -algebra  $A$  with structure map  $\alpha : TA \rightarrow A$ , every arrow  $f : X \rightarrow A$  extends uniquely to a  $T$ -algebra homomorphism  $\bar{f} : TX \rightarrow A$ , which is constructed as  $\bar{f} = \alpha \circ Tf$ :

$$\begin{array}{ccc} X & \xrightarrow{\eta_X} & TX \\ & \searrow f & \nearrow \bar{f} \\ & & A \end{array}$$

See [7, Theorem VI.2.1]. In general, there can be many *arrows*  $\bar{f}$  extending  $f$  along  $\eta_X$ . By 4.3, there is a greatest one, which coincides with the unique  *$T$ -algebra homomorphism* extending  $f$ . In fact, one has that

$$f/\eta_X = \alpha \circ Tf \circ T^*\eta_X \circ \eta_{TX} = \alpha \circ Tf \circ \mu_X \circ \eta_{TX} = \alpha \circ Tf = \bar{f}$$

because  $\alpha = m_A$ . By taking  $A = TY$  and  $f = \eta_Y \circ g$  for  $g : X \rightarrow Y$  arbitrary, recalling that  $Tg : TX \rightarrow TY$  is always an algebra homomorphism, one sees that the functor part of a Kock-Zöberlein monad is uniquely determined by the unit of the monad and by the extension property as  $Tg = (\eta_Y \circ g)/\eta_X$ . In particular, by taking  $g = \text{id}_X$  we conclude that  $\eta_X/\eta_X = \text{id}_{TX}$ .

#### 4.4 *An object is a $T$ -algebra iff it is a retract of a free $T$ -algebra.*

This was first proved by Anders Kock [6]. It is also a corollary of 4.3, using the facts that  $\eta_A$  is a  $T$ -embedding and that an injective object over  $T$ -embeddings is a retract of every object into which it is  $T$ -embedded [3].

## 5 The filter monad is of the Kock-Zöberlein type

$T_0$  topological spaces and continuous maps form a poset-enriched category under the pointwise specialization order. By a simple unfolding of definitions, one sees that the pointwise specialization order is characterized by, for all  $f, g : X \rightarrow Y$ ,

$$f \sqsubseteq g \text{ iff } f^{-1}(V) \subseteq g^{-1}(V) \text{ for every } V \in \Omega Y.$$

### 5.1 *$\mathcal{F}$ is locally monotone.*

Let  $f, g : X \rightarrow Y$  with  $f \sqsubseteq g$ . In order to prove that  $\mathcal{F}f \sqsubseteq \mathcal{F}g$ , let  $\phi \in \mathcal{F}X$  and  $V \in \mathcal{F}f(\phi)$ . This means that  $f^{-1}(V) \in \phi$ . Since  $f^{-1}(V) \subseteq g^{-1}(V)$  and  $\phi$  is upper closed, we have that  $g^{-1}(V) \in \phi$ . But this means that  $V \in \mathcal{F}g(\phi)$ . Therefore  $\mathcal{F}f(\phi) \subseteq \mathcal{F}g(\phi)$ .

## 5.2 $\mathcal{F}$ is a Kock-Zöberlein monad.

By specializing the definitions to the appropriate types, we obtain

$$\eta_{\mathcal{F}X}(\phi) = \{\mathcal{U} \in \Omega\mathcal{F}X \mid \phi \in \mathcal{U}\}, \quad \mathcal{F}\eta_X(\phi) = \{\mathcal{U} \in \Omega\mathcal{F}X \mid \eta_X^{-1}(\mathcal{U}) \in \phi\}.$$

Let  $\mathcal{U} \in \eta_{\mathcal{F}X}(\phi)$ . Then  $\phi \in \mathcal{U}$ . By algebraicity of  $\mathcal{F}X$ , there is  $U \in \phi$  such that already  $\uparrow U \in \mathcal{U}$ . Since  $U = \eta_X^{-1}(\square U) \in \phi$ , we see that  $\square U \in \mathcal{F}\eta_X(\phi)$ . And since  $\square U \subseteq \mathcal{U}$  and filters are upper closed, we conclude that  $\mathcal{U} \in \mathcal{F}\eta_X(\phi)$ . Therefore  $\eta_{\mathcal{F}X}(\phi) \subseteq \mathcal{F}\eta_X(\phi)$ .

## 5.3 The $\mathcal{F}$ -embeddings are precisely the subspace embeddings.

Given a continuous map  $f : X \rightarrow Y$ , its frame map  $f^{-1} : \Omega Y \rightarrow \Omega X$  preserves all joins and hence has a right adjoint  $f_* : \Omega X \rightarrow \Omega Y$ , which sends an open set  $U \in \Omega X$  to the largest open set  $V \in \Omega Y$  such that  $f^{-1}(V) \subseteq U$ . We can thus define a map  $\mathcal{F}^*f : \mathcal{F}Y \rightarrow \mathcal{F}X$  by

$$\mathcal{F}^*f(\gamma) = \{U \in \Omega X \mid f_*(U) \in \gamma\}.$$

The set  $\mathcal{F}^*f(\gamma)$  is a filter because  $f_*$  preserves meets. The function  $\mathcal{F}^*f$  is continuous because one easily computes  $(\mathcal{F}^*f)^{-1}(\square U) = \square f_*(U)$ . That this produces a left adjoint to  $\mathcal{F}f$  is verified as follows:

$$\begin{aligned} \mathcal{F}f(\mathcal{F}^*f(\gamma)) &= \{V \in \Omega Y \mid f^{-1}(V) \in \mathcal{F}^*f(\gamma)\} \\ &= \{V \in \Omega Y \mid f_*(f^{-1}(V)) \in \gamma\} \supseteq \gamma, \end{aligned}$$

because if  $V \in \gamma$  then  $f_*(f^{-1}(V)) \in \gamma$  as  $V \subseteq f_*(f^{-1}(V))$ . Similarly,

$$\begin{aligned} \mathcal{F}^*f(\mathcal{F}f(\phi)) &= \{U \in \Omega X \mid f_*(U) \in \mathcal{F}f(\phi)\} \\ &= \{U \in \Omega X \mid f^{-1}(f_*(U)) \in \phi\} \subseteq \phi. \end{aligned}$$

Reflectiveness means that  $\mathcal{F}^*f \circ \mathcal{F}f(\phi) = \phi$ . So we have to check that the equation  $f^{-1}(f_*(U)) = U$  holds iff  $f$  is an embedding. But this equation is equivalent to saying that  $f^{-1}$  is surjective, which is in turn equivalent to saying that  $f$  is a subspace embedding.

**Note** *The components of the unit are order-monic.*

Assume that  $\eta_X(x) \sqsubseteq \eta_X(y)$  and let  $U$  be an open neighborhood of  $x$ . Then  $U \in \eta_X(x)$  and hence  $U \in \eta_X(y)$  by the assumption, which means that  $y \in U$ . Therefore  $x \sqsubseteq y$ .

**Corollary**  *$\mathcal{F}$  is order-faithful and subspace embeddings are order-monic.*

By virtue of Note 4.1.

## 6 New proofs of old theorems

### 6.1 Continuous lattices

We take the characterization [8, Proposition 2.4] as our definition: A complete lattice  $D$  is *continuous* if every  $d \in D$  is the “lim inf” of its filter of Scott open neighborhoods, in the sense that  $d = \bigsqcup\{\bigsqcap U \mid d \in U\}$ , where  $U$  ranges over Scott open sets.

**6.1.1** *The algebras of the filter monad are the continuous lattices endowed with the Scott topology.*

*Moreover, the structure map  $m_D : \mathcal{F}D \rightarrow D$  of an algebra with underlying space  $D$  is given by*

$$m_D(\phi) = \bigsqcup\{\bigsqcap U \mid U \in \phi\}.$$

We know by 2.1 that the free algebra  $\mathcal{F}X$  is an algebraic lattice endowed with the Scott topology, and by 4.4 that every algebra is a retract of a free algebra. Therefore the algebras are continuous lattices. Conversely, let  $D$  be a continuous lattice endowed with the Scott topology. It is clear that  $m_D : \mathcal{F}D \rightarrow D$  is monotone. We know that  $\mathcal{F}D$  is an algebraic lattice with a basis consisting of principal filters and that every filter is the directed join of the principal filters generated by its members. Thus, in order to establish continuity of  $m_D$ , it is enough to show that  $m_D(\phi) = \bigsqcup\{m_D(\uparrow U) \mid U \in \phi\}$ . But this is immediate because  $m_D(\uparrow U) = \bigsqcap U$ . By §3.1, one has that  $m_D$  is a structure map iff  $\eta_D \circ m_D \sqsubseteq \text{id}_{\mathcal{F}D}$  and  $m_D \circ \eta_D = \text{id}_D$ . The equation holds precisely because  $D$  is continuous. In order to establish the inequality, first notice that  $\eta_D \circ m_D(\phi) = \{U \in \Omega D \mid \bigsqcup\{\bigsqcap U' \mid U' \in \phi\} \in U\}$ . Let  $U \in \eta_D \circ m_D(\phi)$ . Then  $\bigsqcap U' \in U$  for some  $U' \in \phi$  because  $\{\bigsqcap U' \mid U' \in \phi\}$  is directed and  $U$  is Scott open. Hence  $U \in \phi$  because  $U' \subseteq U$ . Therefore  $\eta_D \circ m_D(\phi) \subseteq \phi$ .

**Note** Alan Day also proved that if  $D$  and  $E$  are algebras then a continuous function  $f : D \rightarrow E$  is an algebra homomorphism iff it preserves all meets.

**6.1.2** *The injective spaces over subspace embeddings are the continuous lattices endowed with the Scott topology.*

*Moreover, if  $f : X \rightarrow D$  is a continuous map into a continuous lattice and  $j : X \rightarrow Y$  is a subspace embedding, then  $f$  has a greatest extension along  $j$ , given by*

$$f/j(y) = \bigsqcup\{\bigsqcap U \mid y \in j_*(f^{-1}(U))\}.$$

The  $\mathcal{F}$ -embeddings are the subspace embeddings by 5.3. Therefore the result follows from 4.3, which says that the injective objects over the  $T$ -embeddings of a Kock-Zöberlein monad  $T$  are the  $T$ -algebras. The above formula is a special case of the general formula  $f/j = m_A \circ Tf \circ T^*j \circ \eta_Y$  of 4.3.

**Note** Scott's extension formula produces the map

$$y \mapsto \bigsqcup \left\{ \prod f(j^{-1}(V)) \mid y \in V \right\},$$

which is equivalent to  $f/j$  as defined above, simply because Scott proved that it is the greatest extension along  $j$ .

## 6.2 Continuous Scott domains

The definition of the filter monad still makes sense if the improper filter (the principal filter generated by the empty set—a top element) is ruled out. The resulting monad is referred to as the **proper filter monad** and is denoted by  $\mathcal{F}^+$ . The previous results remain true with the following amendments.

By a **continuous Scott domain** we mean a poset with directed joins and non-empty meets (or, equivalently, bounded joins), subject to the approximation axiom of the definition of a continuous lattice. In the algebraic case one uses the terminology **Scott domain**.

**6.2.1**  $\mathcal{F}^+X$  is a Scott domain endowed with the Scott topology.

**6.2.2** The  $\mathcal{F}^+$ -embeddings are precisely the dense subspace embeddings.

Because a map  $f : X \rightarrow Y$  is dense iff  $f_*(\emptyset) = \emptyset$ , and this is the condition for  $\mathcal{F}^*f(\phi)$  as defined in 5.3 being different from  $\uparrow\emptyset$  for all  $\phi$  and hence  $\mathcal{F}^*f$  being well-defined.

**6.2.3** The algebras of the proper filter monad are the continuous Scott domains endowed with the Scott topology. The homomorphisms are the meet-preserving continuous maps.

**6.2.4** The injective spaces over dense subspace embeddings are the continuous Scott domains endowed with the Scott topology.

**Acknowledgements** I benefited from conversations with Thomas Streicher and Philipp Sünderhauf. I am grateful to Reinhold Heckmann, Anders Kock and Thomas Streicher for (independently) pointing out that a claim made in a previous version of this paper was incorrect. The diagrams were drawn with Paul Taylor's package.

## References

- [1] S. Abramsky and A. Jung. Domain theory. In S. Abramsky, D.M. Gabbay, and T.S.E Maibaum, editors, *Handbook of Logic in Computer Science*, volume 3, pages 1–168. Clarendon Press, Oxford, 1994.
- [2] A. Day. Filter monads, continuous lattices and closure systems. *Canadian Journal of Mathematics*, XXVII(1):50–59, 1975.
- [3] M.H. Escardó. Properly injective spaces and function spaces. *Topology and Its Applications*. To appear (available at <http://www.dcs.ed.ac.uk/home/mhe>).
- [4] M.H. Escardó and R. Flagg. Injective spaces which are the algebras of Kock-Zöberlein monads. In preparation, 1997.
- [5] G. Gierz, K.H. Hofmann, K. Keimel, J.D. Lawson, M. Mislove, and D.S. Scott. *A Compendium of Continuous Lattices*. Springer-Verlag, 1980.
- [6] A. Kock. Monads for which structures are adjoint to units (version 3). *Journal of Pure and Applied Algebra*, 104:41–59, 1995.
- [7] S. Mac Lane. *Categories for the Working Mathematician*. Springer-Verlag, 1971.
- [8] D. S. Scott. Continuous lattices. In F.W. Lawvere, editor, *Toposes, Algebraic Geometry and Logic*, volume 274 of *Lectures Notes in Mathematics*, pages 97–136. Springer-Verlag, 1972.
- [9] M.B. Smyth. Topology. In S. Abramsky, D. M. Gabbay, and T.S.E. Maibaum, editors, *Handbook of Logic in Computer Science*, volume 1, pages 641–761. Clarendon Press, Oxford, 1992.
- [10] S. Vickers. Locales are not pointless. In C. Hankin, I. Mackie, and R. Nagarajan, editors, *Theory and Formal Methods 1994: Proceedings of the Second Imperial College Department of Computing Workshop on Theory and Formal Methods*, Mller Centre, Cambridge, 11–14 September 1994. IC Press. 1995.
- [11] O. Wyler. Algebraic theories for continuous semilattices. *Archive for Rational Mechanics and Analysis*, 90(2):99–113, 1985.